



11 OCT

**EASTERN UNIVERSITY, SRI LANKA**  
**DEPARTMENT OF MATHEMATICS**  
**FIRST EXAMINATION IN SCIENCE - 2011/2012**  
**FIRST SEMESTER (Jan./Feb., 2014)**  
**AM 106 - TENSOR CALCULUS**  
**( Proper & Repeat )**

Time : One hour

Answer all questions

1. (a) Define the following terms:

- i. covariant tensor,
- ii. contravariant tensor.

(b) Write the transformation equation for the following tensors:

- i.  $A_k^{pt}$ ,
- ii.  $B_{tk}^{pqr}$ ,
- iii.  $D_{ptk}^m$ .

(c) Show that the contraction of the outer product of the tensors  $A^p$  and  $B_q$  is an invariant.

(d) The covariant components of a tensor of rank one in rectangular coordinate system are  $2x - z$ ,  $x^2y$ ,  $yz$ . Find its covariant components in spherical coordinate  $(r, \theta, \phi)$ .

- (a) i. Define the Christoffel's symbols of the first and second kind.  
ii. Determine the Christoffel's symbols of the second kind for the metric

$$ds^2 = a^2 d\theta^2 + a^2 \sin^2 \theta d\phi^2$$

where  $a$  is a constant, and find the corresponding differential equation for geodesic.

- (b) i. Write down the covariant derivative of the tensor  $A_{jk}^i$ .

- ii. With the usual notation, prove that

$$\frac{\partial g_{pq}}{\partial x^m} = [pm, q] + [qm, p].$$

Hence deduce that the covariant derivative of a metric tensor  $g_{jk}$  is zero.

- iii. Using the covariant derivative of a metric tensor, prove that

$$\Gamma_{ca}^e = \frac{1}{2} g^{eb} [\partial_c(g_{ab}) + \partial_a(g_{cb}) - \partial_b(g_{ca})], \quad \text{where } \partial_i = \frac{\partial}{\partial x^i}.$$