



EASTERN UNIVERSITY, SRI LANKA
DEPARTMENT OF MATHEMATICS
FIRST EXAMINATION IN SCIENCE - 2011/2012
FIRST SEMESTER (Jan./Feb., 2014)
MT 106 - TENSOR CALCULUS
(Re-Repeat)

Answer all questions

Time : One hour

1. (a) Define the following terms:

- i. covariant tensor,
- ii. contravariant tensor.

(b) Write the transformation equation for the following tensors:

- i. A_k^p ,
- ii. B_{lk}^{pq} ,
- iii. D_{ptk}^m .

(c) Show that the contraction of the outer product of the tensors A^p and B_q is an invariant.

(d) The covariant components of a tensor of rank one in rectangular coordinate system are $2x - z$, x^2y , yz . Find its covariant components in spherical coordinate (r, θ, ϕ) .

2. (a) i. Define the Christoffel's symbols of the first and second kind.
ii. Determine the Christoffel's symbols of the second kind for the metric

$$ds^2 = a^2 d\theta^2 + a^2 \sin^2 \theta d\phi^2$$

where a is a constant, and find the corresponding differential equation for geodesic.

- (b) i. Write down the covariant derivative of the tensor A_{jk}^i .

- ii. With the usual notation, prove that

$$\frac{\partial g_{pq}}{\partial x^m} = [pm, q] + [qm, p].$$

Hence deduce that the covariant derivative of a metric tensor g_{jk} is zero.

- iii. Using the covariant derivative of a metric tensor, prove that

$$\Gamma_{ca}^e = \frac{1}{2} g^{eb} [\partial_c(g_{ab}) + \partial_a(g_{cb}) - \partial_b(g_{ca})], \quad \text{where } \partial_i = \frac{\partial}{\partial x^i}.$$